


-198-

hemangiomas, hepatic adenoma, hepatic adenomatosis,  
hepatocellular carcinoma, insulinoma, intraepithelial  
neoplasia, interepithelial squamous cell neoplasia,  
invasive squamous cell carcinoma, large cell carcinoma,  
5 leiomyosarcoma, lentigo maligna melanomas, malignant  
melanoma, malignant mesothelial tumors, medulloblastoma,  
medulloepithelioma, melanoma, meningeal, mesothelial,  
metastatic carcinoma, mucoepidermoid carcinoma,  
neuroblastoma, neuroepithelial adenocarcinoma nodular  
10 melanoma, oat cell carcinoma, oligodendroglial,  
osteosarcoma, pancreatic polypeptide, papillary serous  
adenocarcinoma, pineal cell, pituitary tumors,  
plasmacytoma, pseudosarcoma, pulmonary blastoma, renal  
cell carcinoma, retinoblastoma, rhabdomyosarcoma,  
15 sarcoma, serous carcinoma, small cell carcinoma, soft  
tissue carcinomas, somatostatin-secreting tumor,  
squamous carcinoma, squamous cell carcinoma,  
submesothelial, superficial spreading melanoma,  
undifferentiated carcinoma, uveal melanoma, verrucous  
20 carcinoma, vipoma, well differentiated carcinoma, and  
Wilm's tumor.

46. The method of Claim 44 wherein the  
cyclooxygenase-2 inhibitor is selected from compounds,  
and their pharmaceutically acceptable salts thereof, of  
25 the group consisting of:

-199-

1)



JTE-522, 4-(4-cyclohexyl-2-methyloxazol-5-yl)-  
2-fluorobenzenesulfonamide,

5

2)

5-chloro-3-(4-(methylsulfonyl)phenyl)-2-  
(methyl-5-pyridinyl)pyridine,


3)

2-(3,5-difluorophenyl)-3-4-

10

(methylsulfonyl)phenyl)-2-cyclopenten-1-one,


4)



4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-  
pyrazol-1-yl]-benzenesulfonamide,

-200-


5)



rofecoxib, 4-(4-(methylsulfonyl)phenyl)-3-phenyl-2(5H)-furanone,

5

6)




4-(5-methyl-3-phenylisoxazol-4-yl)benzenesulfonamide,

7) N-[[4-(5-methyl-3-phenylisoxazol-4-yl)phenyl]sulfonyl]propanamide,

10

8)



4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazole-1-yl]benzenesulfonamide,